

Package: CauchyCP (via r-universe)

August 26, 2024

Type Package

Title Powerful Test for Survival Data under Non-Proportional Hazards

Version 0.1.1

Author Hong Zhang

Maintainer Hong Zhang <hzhang@wpi.edu>

Description An omnibus test of change-point Cox regression models to improve the statistical power of detecting signals of non-proportional hazards patterns. The technical details can be found in Hong Zhang, Qing Li, Devan Mehrotra and Judong Shen (2021) <[arXiv:2101.00059](https://arxiv.org/abs/2101.00059)>. Extensive simulation studies demonstrate that, compared to existing tests under non-proportional hazards, the proposed CauchyCP test 1) controls the type I error better at small alpha levels; 2) increases the power of detecting time-varying effects; and 3) is more computationally efficient.

License GPL-2

Imports stats, survival

Encoding UTF-8

LazyData true

RoxygenNote 6.1.0

NeedsCompilation no

Date/Publication 2022-08-12 06:10:06 UTC

Repository <https://wpihongzhang.r-universe.dev>

RemoteUrl <https://github.com/cran/CauchyCP>

RemoteRef HEAD

RemoteSha 9f104f8b81f3c99c0e676f3e66f529bcbb91f8d7

Contents

CauchyCP	2
gast	3

Index**4**

CauchyCP	<i>A robust test under non-proportional hazards using Cauchy combination of change-point Cox regressions.</i>
----------	---

Description

A robust test under non-proportional hazards using Cauchy combination of change-point Cox regressions.

Usage

```
CauchyCP(time, status, x, covar = rep(1, length(time)),
          cutpoints = c(0, quantile(time[status == 1])[2:4]))
```

Arguments

time	- Follow up time for right censored data.
status	- The event status indicator, 0=censored, 1=event.
x	- The variable of interest, e.g. a treatment indicator.
covar	- The matrix of covariates. If no covariates, a vector of ones should be used (default).
cutpoints	- The pre-specified change-points. The default choice is a vector of 0th, 25th, 50th and 75th percentiles of the event time.

Value

1. A matrix of estimated hazard ratios before and after the change-points. 2. the vector of p-values corresponding to the change-points. 3. a final p-value.

References

Hong Zhang, Qing Li, Devan Mehrotra and Judong Shen. "CauchyCP: a powerful test under non-proportional hazards using Cauchy combination of change-point Cox regressions", arXiv:2101.00059.

Examples

```
data(gast)
CauchyCP(time=gast$time, status=gast$status, x=gast$trt)
```

gast

Example 1: gastric carcinoma trial data

Description

A two-arm gastric carcinoma clinical trial: ninety patients with locally advanced, non-resectable gastric carcinoma received either chemotherapy alone (N = 45) or chemotherapy plus radiation (N = 45).

Usage

gast

Format

A data frame with 90 rows and 3 variables:

trt treatment indicator, 1=chemotherapy + radiation, 0=chemotherapy alone.

status event indicator, 1=death, 0=censored.

time follow up time, in days

Source

K. R. Hess, Assessing time-by-covariate interactions in proportional hazards regression models using cubic spline functions, *Statistics in medicine* 13 (10) (1994) 1045–1062.

Index

* **datasets**
gast, 3

CauchyCP, 2

gast, 3